

Analytical Blowup Solutions to the Pressureless Navier-Stokes-Poisson Equations with Density-dependent Viscosity in R^N

YUEN MANWAI*

*Department of Applied Mathematics,
The Hong Kong Polytechnic University,
Hung Hom, Kowloon, Hong Kong*

Revised 24-Apr-2009

Abstract

We study the N-dimensional pressureless Navier-Stokes-Poisson equations with density-dependent viscosity. With the extension of the blowup solutions for the Euler-Poisson equations, the analytical blowup solutions, in radial symmetry, in R^N ($N \geq 2$) are constructed.

1 Introduction

The evolution of a self-gravitating fluid can be formulated by the Navier-Stokes-Poisson equations of the following form:

$$\begin{cases} \rho_t + \nabla \cdot (\rho \vec{u}) = 0, \\ (\rho \vec{u})_t + \nabla \cdot (\rho \vec{u} \otimes \vec{u}) + \nabla P = -\rho \nabla \Phi + \text{vis}(\rho, \vec{u}), \\ \Delta \Phi(t, x) = \alpha(N)\rho, \end{cases} \quad (1)$$

where $\alpha(N)$ is a constant related to the unit ball in R^N : $\alpha(1) = 2$; $\alpha(2) = 2\pi$ and For $N \geq 3$,

$$\alpha(N) = N(N-2)V(N) = N(N-2)\frac{\pi^{N/2}}{\Gamma(N/2+1)}, \quad (2)$$

where $V(N)$ is the volume of the unit ball in R^N and Γ is a Gamma function. And as usual, $\rho = \rho(t, x)$ and $\vec{u} = \vec{u}(t, x) \in \mathbf{R}^N$ are the density, the velocity respectively. $P = P(\rho)$ is the pressure. The evolution of the cosmology can be modelled by the dust distribution without pressure term. That describes the stellar systems of collisionless, gravitational N-body systems.

*E-mail address: nevetsyuen@hotmail.com

In the above system, the self-gravitational potential field $\Phi = \Phi(t, x)$ is determined by the density ρ through the Poisson equation.

And $vis(\rho, \vec{u})$ is the viscosity function:

$$vis(\rho, \vec{u}) := \nabla(\mu(\rho)) \nabla \bullet \vec{u}. \quad (3)$$

In this article, we seek the radial solutions

$$\rho(t, x) = \rho(t, r) \text{ and } \vec{u} = \frac{\vec{x}}{r} V(t, r) := \frac{\vec{x}}{r} V, \quad (4)$$

with $r = \left(\sum_{i=1}^N x_i^2 \right)^{1/2}$. In particular, we can deduce the viscosity function (3) to be

$$\nabla(\mu(\rho)) \nabla \bullet \vec{u} = \nabla \left[\mu(\rho) \left(\left(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_N} \right) \cdot \left(\frac{x_1}{r}, \frac{x_2}{r}, \dots, \frac{x_N}{r} \right) V \right) \right] \quad (5)$$

$$= \nabla \left[\mu(\rho) \sum_{k=1}^N \frac{\partial}{\partial x_k} \left(\frac{x_k}{r} V \right) \right] \quad (6)$$

$$= \nabla \left[\mu(\rho) \sum_{k=1}^n \left(\frac{r^2 - x_k^2}{r^3} V + \frac{x_k}{r} V_r \frac{x_k}{r} \right) \right] \quad (7)$$

$$= \nabla \left[\mu(\rho) \left(\frac{(N-1)r^2}{r^3} V + V_r \right) \right] \quad (8)$$

$$= \nabla \left[\mu(\rho) \left(\frac{N-1}{r} V + V_r \right) \right] \quad (9)$$

$$= \nabla \mu(\rho) \left(\frac{N-1}{r} V + V_r \right) + \mu(\rho) \nabla \left(\frac{N-1}{r} V + V_r \right) \quad (10)$$

$$= \frac{\partial}{\partial x_i} \mu_r(\rho) \left(\frac{N-1}{r} V + V_r \right) + \mu(\rho) \frac{\partial}{\partial x_i} \left(\frac{N-1}{r} V + V_r \right) \quad (11)$$

$$= \frac{\partial r}{\partial x_i} \mu_r(\rho) \left(\frac{N-1}{r} V + V_r \right) + \mu(\rho) \left(- \left(\frac{N-1}{r^2} \right) \frac{\partial r}{\partial x_i} V + \frac{N-1}{r} \frac{\partial r}{\partial x_i} V_r + \frac{\partial r}{\partial x_i} V_{rr} \right) \quad (12)$$

$$= \frac{x_i}{r} \mu_r(\rho) \left(\frac{N-1}{r} V + V_r \right) + \mu(\rho) \left(- \left(\frac{N-1}{r^2} \right) \frac{x_i}{r} V + \frac{N-1}{r} \frac{x_i}{r} V_r + \frac{x_i}{r} V_{rr} \right) \quad (13)$$

$$= \frac{x_i}{r} \left\{ \mu_r(\rho) \left(\frac{N-1}{r} V + V_r \right) + \mu(\rho) \left[- \left(\frac{N-1}{r^2} \right) V + \frac{N-1}{r} V_r + V_{rr} \right] \right\} \quad (14)$$

The equations (3) can be converted to the radial solutions to have

$$\rho [u_t + (u \cdot \nabla) u] + \nabla P(\rho) + \rho \nabla \Phi \quad (15)$$

$$= \rho \left((u_i)_t + \sum_{k=1}^n u_k \frac{\partial u_i}{\partial x_k} \right) + \frac{\partial P(\rho)}{\partial x_i} + \rho \frac{\partial \Phi(\rho)}{\partial x_i}, \text{ for } i = 1, 2, \dots, N \quad (16)$$

$$= \rho \left((u_i)_t + u_i \frac{\partial u_i}{\partial x_i} + \sum_{k \neq i} u_k \frac{\partial u_i}{\partial x_k} \right) + \frac{\partial P(\rho)}{\partial x_i} + \rho \frac{\partial \Phi(\rho)}{\partial x_i} \quad (17)$$

$$= \rho \left[\frac{\partial}{\partial t} \left(\frac{x_i}{r} V \right) + \frac{x_i}{r} V \frac{\partial}{\partial x_i} \left(\frac{x_i}{r} V \right) + \sum_{k \neq i} \frac{x_k}{r} V \frac{\partial}{\partial x_k} \left(\frac{x_i}{r} V \right) \right] + \left(\frac{\partial P(\rho)}{\partial r} + \rho \frac{\partial \Phi(\rho)}{\partial r} \right) \frac{\partial r}{\partial x_i} \quad (18)$$

$$= \rho \left[\frac{x_i}{r} V_t + \frac{x_i}{r} V \left(\frac{r^2 - x_i^2}{r^3} V + \frac{x_i}{r} V_r \frac{x_i}{r} \right) \right] + \sum_{k \neq i} \frac{x_k}{r} V \left(\frac{-x_i x_k}{r^3} V + \frac{x_i x_k}{r^2} V_r \right) + [P_r(\rho) + \rho \Phi_r(\rho)] \frac{x_i}{r} \quad (19)$$

$$= \frac{x_i}{r} \left\{ \rho \left[V_t + \left(\frac{r^2 - x_i^2}{r^3} - \sum_{k \neq i} \frac{x_k^2}{r^3} \right) V^2 + \left(\frac{x_i^2}{r^3} + \sum_{k \neq i} \frac{x_k^2}{r^3} \right) V V_r \right] + P_r(\rho) + \rho \Phi_r(\rho) \right\} \quad (20)$$

$$= \frac{x_i}{r} [\rho (V_t + V V_r) + P_r(\rho) + \rho \Phi_r(\rho)]. \quad (21)$$

Therefore, the Euler-Poisson equations in radial symmetry can be written in the following form

$$\begin{cases} \rho_t + V \rho_r + \rho V_r + \frac{N-1}{r} \rho V = 0, \\ \rho (V_t + V V_r) + P_r(\rho) + \rho \Phi_r(\rho) = \mu_r(\rho) \left(\frac{N-1}{r} V + V_r \right) + \mu(\rho) \left[- \left(\frac{N-1}{r^2} \right) V + \frac{N-1}{r} V_r + V_{rr} \right]. \end{cases} \quad (22)$$

Here we under a common assumption for:

$$\mu(\rho) := \kappa \rho^\theta, \quad (23)$$

and κ and $\theta \geq 0$ are the constants. For the study of this kind of the above system, the readers may refer [7], [9], [10], [12] and [14]. In particular, when $\theta = 0$, it returns the expression for the V -dependent only viscosity function:

$$\begin{cases} \rho_t + V \rho_r + \rho V_r + \frac{N-1}{r} \rho V = 0, \\ \rho (V_t + V V_r) + P_r(\rho) + \rho \Phi_r(\rho) = \mu(\rho) \left[V_{rr} + \frac{N-1}{r} V_r - \left(\frac{N-1}{r^2} \right) V \right], \end{cases} \quad (24)$$

which are the usual form Navier-Stokes-Poisson equations. The equations (1)₁ and (1)₂ ($vis(\rho, u) \neq 0$) are the compressible Navier-Stokes equations with forcing term. The equation (1)₃ is the Poisson equation through which the gravitational potential is determined by the density distribution of the density itself. Thus, we call the system (1) the Navier-Stokes-Poisson equations.

Here, if the $vis(\rho, u) = 0$, the system is called the Euler-Poisson equations. In this case, the equations can be viewed as a perfect gas model. For $N = 3$, (1) is a classical (nonrelativistic) description of a galaxy, in astrophysics. See [2], [8] for a detail about the system.

$P = P(\rho)$ is the pressure. The γ -law can be applied on the pressure $P(\rho)$, i.e.

$$P(\rho) = K \rho^\gamma := \frac{\rho^\gamma}{\gamma}, \quad (25)$$

which is a commonly the hypothesis. The constant $\gamma = c_P/c_v \geq 1$, where c_P, c_v are the specific heats per unit mass under constant pressure and constant volume respectively, is the ratio of the specific heats, that is, the adiabatic exponent in (25). With $K = 0$, we call that the system is pressureless.

For the 3-dimensional case, we are interested in the hydrostatic equilibrium specified by $u = 0$. According to [2], the ratio between the core density $\rho(0)$ and the mean density $\bar{\rho}$ for $6/5 < \gamma < 2$ is given by

$$\frac{\bar{\rho}}{\rho(0)} = \left(\frac{-3}{z} \dot{y}(z) \right)_{z=z_0} \quad (26)$$

where y is the solution of the Lane-Emden equation with $n = 1/(\gamma - 1)$,

$$\ddot{y}(z) + \frac{2}{z} \dot{y}(z) + y(z)^n = 0, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \quad n = \frac{1}{\gamma - 1}, \quad (27)$$

and z_0 is the first zero of $y(z_0) = 0$. We can solve the Lane-Emden equation analytically for

$$y_{anal}(z) \doteq \begin{cases} 1 - \frac{1}{6}z^2, & n = 0; \\ \frac{\sin z}{z}, & n = 1; \\ \frac{1}{\sqrt{1 + z^2/3}}, & n = 5, \end{cases} \quad (28)$$

and for the other values, only numerical values can be obtained. It can be shown that for $n < 5$, the radius of polytropic models is finite; for $n \geq 5$, the radius is infinite.

For the isothermal case ($\gamma = 1$), the corresponding stationary solution is the Liouville equation,

$$\ddot{y}(z) + \frac{2}{z} \dot{y}(z) + \frac{\alpha(3)}{K} e^{y(z)} = 0. \quad (29)$$

Gambin [4] and Bezard [1] obtained the existence results about the explicitly stationary solution ($u = 0$) for $\gamma = 6/5$ in Euler-Poisson equations:

$$\rho = \left(\frac{3KA^2}{2\pi} \right)^{5/4} (1 + A^2 r^2)^{-5/2}, \quad (30)$$

where A is constant.

In the following, we always seek solutions in radial symmetry. Thus, the Poisson equation (1)₃ is transformed to

$$r^{N-1} \Phi_{rr}(t, x) + (N-1) r^{N-2} \Phi_r = \alpha(N) \rho r^{N-1}, \quad (31)$$

$$\Phi_r = \frac{\alpha(N)}{r^{N-1}} \int_0^r \rho(t, s) s^{N-1} ds. \quad (32)$$

In this paper, we concern about blowup solutions for the N -dimensional pressureless Navier-Stokes-Poisson equations with the density-dependent viscosity. And our aim is to construct a family of such blowup solutions.

Historically in astrophysics, Goldreich and Weber [5] constructed the analytical blowup solutions (collapsing) of the 3-dimensional Euler-Poisson equations for $\gamma = 4/3$ for the non-rotating

gas spheres. After that, Makino [8] obtained the rigorously mathematical proof of the existence of such kind of blowup solutions. And in [3], Deng, Xiang and Yang extended the above blowup solutions in R^N ($N \geq 4$). Recently, Yuen obtained the blowup solutions in R^2 with $\gamma = 1$ in [14]. The family of the analytical solutions are rewritten as

For $N \geq 3$ and $\gamma = (2N - 2)/N$, in [13]

$$\left\{ \begin{array}{l} \rho(t, r) = \begin{cases} \frac{1}{a(t)^N} y\left(\frac{r}{a(t)}\right)^{N/(N-2)}, & \text{for } r < a(t)Z_\mu; \\ 0, & \text{for } a(t)Z_\mu \leq r. \end{cases}, \quad V(t, r) = \frac{\dot{a}(t)}{a(t)} r, \\ \ddot{a}(t) = -\frac{\lambda}{a(t)^{N-1}}, \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1, \\ \ddot{y}(z) + \frac{N-1}{z} \dot{y}(z) + \frac{\alpha(N)}{(2N-2)K} y(z)^{N/(N-2)} = \mu, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \end{array} \right. \quad (33)$$

where $\mu = [N(N-2)\lambda]/(2N-2)K$ and the finite Z_μ is the first zero of $y(z)$;

For $N = 2$ and $\gamma = 1$, in [14]

$$\left\{ \begin{array}{l} \rho(t, r) = \frac{1}{a(t)^2} e^{y(r/a(t))}, \quad V(t, r) = \frac{\dot{a}(t)}{a(t)} r; \\ \ddot{a}(t) = -\frac{\lambda}{a(t)}, \quad a(0) = a_0 > 0, \quad \dot{a}(0) = a_1; \\ \ddot{y}(z) + \frac{1}{z} \dot{y}(z) + \frac{\alpha(2)}{K} e^{y(z)} = \mu, \quad y(0) = \alpha, \quad \dot{y}(0) = 0, \end{array} \right. \quad (34)$$

where $K > 0$, $\mu = 2\lambda/K$ with a sufficiently small λ and α are constants.

For the construction of special analytical solutions to the Navier-Stokes equations in R^N without force term, readers may refer Yuen's recent results in [15].

In this article, the analytical blowup solutions are constructed in the pressureless Euler-Poisson equations with density-dependent viscosity in R^N in radial symmetry:

$$\left\{ \begin{array}{l} \rho_t + V\rho_r + \rho V_r + \frac{N-1}{r} \rho V = 0, \\ \rho(V_t + VV_r) = -\frac{\alpha(N)\rho}{r^{N-1}} \int_0^r \rho(t, s) s^{N-1} ds + [\kappa \rho^\theta]_r \left(\frac{N-1}{r} V + V_r \right) + (\kappa \rho^\theta)(V_{rr} + \frac{N-1}{r} V_r + \frac{N-1}{r^2} V), \end{array} \right. \quad (35)$$

in the form of the following theorem.

Theorem 1 *For the N -dimensional pressureless Navier-Stokes-Poisson equations in radial symmetry, (35), there exists a family of solutions:*

for $N \geq 2$ and $N \neq 3$, with $\theta = (2N - 3)/N$, we have,

$$\left\{ \begin{array}{l} \rho(t, r) = \begin{cases} \frac{1}{(T - Ct)^N} y\left(\frac{r}{T - Ct}\right)^{\frac{N}{N-3}}, & \text{for } \frac{r}{T - Ct} < Z_0; \\ 0, & \text{for } Z_0 \leq \frac{r}{T - Ct}. \end{cases}, \quad V(t, r) = \frac{-C}{T - Ct} r; \\ \ddot{y}(z) + \frac{N-1}{z} \dot{y}(z) + \frac{\alpha(N)(N-3)}{(2N-3)CN\kappa} y(z)^{N/(N-3)} = 0, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0 \end{array} \right.; \quad (36)$$

for $N = 3$, with $\theta = 1$, we have,

$$\left\{ \begin{array}{l} \rho(t, r) = \frac{1}{(T - Ct)^3} e^{y(r/(T - Ct))}, \quad V(t, r) = \frac{-C}{T - Ct} r; \\ \ddot{y}(z) + \frac{2}{z} \dot{y}(z) + \frac{4\pi}{CN\kappa} e^{y(z)} = 0, \quad y(0) = \alpha, \quad \dot{y}(0) = 0, \end{array} \right. \quad (37)$$

where $T > 0$, $\kappa > 0$, $C \neq 0$ and α are constants, and the finite Z_0 is the first zero of $y(z)$. In particular, for $C > 0$, the solutions blow up in the finite time T/C .

2 Separable Blowup Solutions

Before presenting the proof of Theorem 1, we prepare some lemmas. First, we obtain the solutions for the continuity equation of mass in radial symmetry (35)₁.

Lemma 2 *For the N -dimensional conservation of mass in radial symmetry*

$$\rho_t + V\rho_r + \rho V_r + \frac{N-1}{r}\rho V = 0, \quad (38)$$

there exist solutions,

$$\rho(t, r) = \frac{f(r/(T - Ct))}{(T - Ct)^N}, \quad V(t, r) = \frac{-C}{T - Ct}r, \quad (39)$$

where $f \geq 0 \in C^1$, T and C are positive constants.

Proof. We just plug (39) into (38). Then

$$\rho_t + V\rho_r + \rho V_r + \frac{N-1}{r}\rho V \quad (40)$$

$$= \frac{(-N)(-C)f(r/(T - Ct))}{(T - Ct)^{N+1}} - \frac{(-C)r\dot{f}(r/(T - Ct))}{(T - Ct)^{N+2}} \quad (41)$$

$$+ \frac{(-C)r}{T - Ct} \frac{\dot{f}(r/(T - Ct))}{(T - Ct)^{N+1}} + \frac{f(r/(T - Ct))}{(T - Ct)^N} \frac{(-C)}{T - Ct} + \frac{N-1}{r} \frac{f(r/(T - Ct))}{(T - Ct)^N} \frac{(-C)}{T - Ct}r \quad (42)$$

$$= 0. \quad (43)$$

The proof is completed. ■

Besides, we need the lemma for stating the property of the function $y(z)$ of the analytical solutions (36) and (37). We need the lemma for stating the property of the function $y(z)$. In particular, the solutions (36) in 2-dimensional case involve the following lemma. The similar lemma was already given in Lemmas 9 and 10, in [14], by the fixed point theorem. For the completeness of understanding the whole article, the proof is also presented here.

Lemma 3 *For the ordinary differential equation,*

$$\begin{cases} \ddot{y}(z) + \frac{1}{z}\dot{y}(z) - \frac{\sigma}{y(z)^2} = 0 \\ y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \end{cases} \quad (44)$$

where σ is a positive constant,

has a solution $y(z) \in C^2$ and $\lim_{z \rightarrow +\infty} y(z) = \infty$.

Proof. By integrating (44), we have,

$$\dot{y}(z) = \frac{\sigma}{z} \int_0^z \frac{1}{y(s)^2} s ds \geq 0. \quad (45)$$

Thus, for $0 < z < z_0$, $y(z)$ has a uniform lower upper bound

$$y(z) \geq y(0) = \alpha > 0. \quad (46)$$

As we obtained the local existence in Lemma 3, there are two possibilities:

(1) $y(z)$ only exists in some finite interval $[0, z_0]$: (1a) $\lim_{z \rightarrow z_0^-} y(z) = \infty$; (1b) $y(z)$ has an uniformly upper bound, i.e. $y(z) \leq \alpha_0$ for some constant α_0 .

(2) $y(z)$ exists in $[0, +\infty)$: (2a) $\lim_{z \rightarrow +\infty} y(z) = \infty$; (2b) $y(z)$ has an uniformly upper bound, i.e. $y(z) \leq \beta$ for some positive constant β .

We claim that possibility (1) does not exist. We need to reject (1b) first: If the statement (1b) is true, (45) becomes

$$\frac{\sigma z}{2\alpha^2} = \frac{\sigma}{z} \int_0^z \frac{s}{\alpha^2} ds \geq \dot{y}(z). \quad (47)$$

Thus, $\dot{y}(z)$ is bounded in $[0, z_0]$. Therefore, we can use the fixed point theorem again to obtain a large domain of existence, such that $[0, z_0 + \delta]$ for some positive number δ . There is a contradiction. Therefore, (1b) is rejected.

Next, we do not accept (1a) because of the following reason: It is impossible that $\lim_{z \rightarrow z_0^-} y(z) = \infty$, as from (47), $\dot{y}(z)$ has an upper bound in $[0, z_0]$:

$$\frac{\sigma z_0}{2\alpha^2} \geq \dot{y}(z). \quad (48)$$

Thus, (48) becomes,

$$y(z_0) = y(0) + \int_0^{z_0} \dot{y}(s) ds \leq \alpha + \int_0^{z_0} \frac{\sigma z_0}{2\alpha^2} ds = \alpha + \frac{\sigma(z_0)^2}{2\alpha^2} \quad (49)$$

Since $y(z)$ is bounded above in $[0, z_0]$, it contradicts the statement (1a), such that $\lim_{z \rightarrow z_0^-} y(z) = \infty$. So, we can exclude the possibility (1).

We claim that the possibility (2b) doesn't exist. It is because

$$\dot{y}(z) = \frac{\sigma}{z} \int_0^z \frac{s}{y(s)^2} ds \geq \frac{\sigma}{z} \int_0^z \frac{s}{\beta^2} ds = \frac{\sigma z}{2\beta^2}. \quad (50)$$

Then, we have,

$$y(z) \geq \alpha + \frac{\sigma}{4\beta^2} z^2. \quad (51)$$

By letting $z \rightarrow \infty$, (51) turns out to be,

$$y(z) = \infty. \quad (52)$$

Since a contradiction is established, we exclude the possibility (2b). Thus, the equation (44) exists in $[0, +\infty)$ and $\lim_{z \rightarrow +\infty} y(z) = \infty$. This completes the proof. ■

For $N \geq 4$, our blowup solutions depend on the Lane-Emden equation,

$$\begin{cases} \ddot{y}(z) + \frac{N-1}{z} \dot{y}(z) + \sigma y(z)^s = 0, \\ y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \end{cases} \quad (53)$$

where $s > 1$ and σ are positive constants, which is a particular case of the Emden-Fowler equation

$$\ddot{u} + z^{1-n} u^n = 0, \quad (54)$$

with $n > 1$, after the transformation

$$u = \frac{y(z)}{z}. \quad (55)$$

For the existence and uniqueness of the equation (53), the interested readers may refer the survey paper [11].

The proof of Theorem 1 is similar to the ones in [3], [13] and [14]. The main idea is to put the analytical solutions to check that if they satisfy the system (35) only.

Proof of Theorem 1. From Lemma 2, it is clear for that (36) satisfy (35)₁.

For the case of $N \geq 2$ and $N \neq 3$ with $\theta = (2N-3)/N$, we plug the solutions (36) into the momentum equation (35)₂,

$$\rho(V_t + VV_r) + \frac{\alpha(N)\rho}{r^{N-1}} \int_0^r \rho(t, s) s^{N-1} ds - [\kappa \rho^{(2N-3)/N}]_r \left(\frac{N-1}{r} V + V_r \right) - \kappa \rho^{(2N-3)/N} (V_{rr} + \frac{N-1}{r} V_r - \frac{N-1}{r^2} V) \quad (56)$$

$$= \rho \left[\frac{(-C)(-1)(-C)}{(T-Ct)^2} r + \frac{(-C)}{T-Ct} r \cdot \frac{(-C)}{T-Ct} \right] + \frac{\alpha(N)\rho}{r^{N-1}} \int_0^r \frac{y(\frac{s}{T-Ct})^{N/(N-3)}}{(T-Ct)^N} s^{N-1} ds \quad (57)$$

$$- \frac{\partial}{\partial r} N\kappa \left[\frac{1}{(T-Ct)^N} y \left(\frac{r}{T-Ct} \right)^{N/(N-3)} \right]^{(2N-3)/N} \frac{(-C)}{T-Ct} - 0 \quad (58)$$

$$= \frac{\alpha(N)\rho}{r^{N-1}} \int_0^r \frac{y(\frac{s}{T-Ct})^{N/(N-3)}}{(T-Ct)^N} s^{N-1} ds + \frac{(2N-3)CN\kappa}{N-3} \frac{y(\frac{r}{T-Ct})}{(T-Ct)^{N-3}} \frac{y(\frac{r}{T-Ct})^{[N/(N-3)]-1}}{(T-Ct)^{N+2}} \dot{y}\left(\frac{r}{T-Ct}\right) \quad (59)$$

$$= \frac{\alpha(N)\rho}{r^{N-1}} \int_0^r \frac{y(\frac{s}{T-Ct})^{N/(N-3)}}{(T-Ct)^N} s^{N-1} ds + \frac{(2N-3)CN\kappa}{N-3} \frac{\rho}{(T-Ct)^{N-1}} \dot{y}\left(\frac{r}{T-Ct}\right) \quad (60)$$

$$= \frac{\rho}{(T-Ct)^{N-1}} \left[\frac{(2N-3)CN\kappa}{N-3} \dot{y}\left(\frac{r}{T-Ct}\right) + \frac{\alpha(N)}{r^{N-1}(T-Ct)} \int_0^r y\left(\frac{s}{T-Ct}\right)^{N/(N-3)} s^{N-1} ds \right] \quad (61)$$

$$= \frac{\rho}{(T-Ct)^{N-1}} \left[\frac{(2N-3)CN\kappa}{N-3} \dot{y}\left(\frac{r}{T-Ct}\right) + \frac{\alpha(N)}{(\frac{r}{T-Ct})^{N-1}} \int_0^{r/(T-Ct)} y(s)^{N/(N-3)} s^{N-1} ds \right] \quad (62)$$

$$= \frac{\rho}{(T-Ct)^{N-1}} Q\left(\frac{r}{T-Ct}\right). \quad (63)$$

And denote

$$Q\left(\frac{r}{T-Ct}\right) \doteq Q(z) = \frac{(2N-3)CN\kappa}{N-3} \dot{y}(z) + \frac{\alpha(N)}{z^{N-1}} \int_0^z y(s)^{N/(N-3)} s^{N-1} ds. \quad (64)$$

Differentiate $Q(z)$ with respect to z ,

$$\dot{Q}(z) = \frac{(2N-3)CN\kappa}{N-3} \ddot{y}(z) + \alpha(N) \dot{y}(z)^{N/(N-3)} - \frac{(N-1)\alpha(N)}{z^{N-1}} \int_0^z y(s)^{N/(N-3)} s^{N-1} ds \quad (65)$$

$$= -\left(\frac{N-1}{z}\right) \frac{(2N-3)CN\kappa}{N-3} \dot{y}(z) - \frac{(N-1)\alpha(N)}{z} \frac{\alpha(N)}{z^{N-1}} \int_0^z y(s)^{N/(N-3)} s^{N-1} ds \quad (66)$$

$$= \frac{-(N-1)}{z} Q(z), \quad (67)$$

where the above result is due to the fact that we choose the following ordinary differential equation,

$$\begin{cases} \ddot{y}(z) + \frac{N-1}{z} \dot{y}(z) + \frac{\alpha(N)(N-3)}{(2N-3)CN\kappa} y(z)^{N/(N-3)} = 0, \\ y(0) = \alpha > 0, \dot{y}(0) = 0, \end{cases} \quad (68)$$

in Lemma 3 and the well-known results in the Emden-Fowler equation. With $Q(0) = 0$, this implies that $\dot{Q}(z) = 0$. Thus, the momentum equation $(35)_2$ is satisfied.

For the case of $N = 3$ with $\theta = 1$, we plug the solutions (37) into the momentum equation $(35)_2$,

$$\rho(V_t + VV_r) + \frac{4\pi\rho}{r^2} \int_0^r \rho(t, s) s^2 ds - [\mu(\rho)]_r \left(\frac{N-1}{r} V + V_r \right) - \mu(\rho) \left(V_{rr} + \frac{2}{r} V_r - \frac{2}{r^2} V \right) \quad (69)$$

$$= \rho \left[\frac{(-C)(-1)(-C)}{(T-Ct)^2} r + \frac{(-C)}{T-Ct} r \cdot \frac{(-C)}{T-Ct} \right] + \frac{4\pi\rho}{r^2} \int_0^r \frac{e^{y(s/(T-Ct))}}{(T-Ct)^3} s^2 ds \quad (70)$$

$$- (\kappa\rho)_r \frac{N(-C)}{T-Ct} - \mu(\rho) \left(0 + \frac{2}{r} \frac{(-1)}{T-Ct} - \frac{2}{r^2} \frac{(-1)}{T-Ct} r \right) \quad (71)$$

$$= \frac{\kappa e^{y(r/(T-Ct))}}{(T-Ct)^3} \cdot \dot{y}\left(\frac{r}{T-Ct}\right) \cdot \frac{1}{T-Ct} \frac{NC}{T-Ct} + \frac{4\pi\rho}{r^2} \int_0^r \frac{e^{y(s/(T-Ct))}}{(T-Ct)^3} s^2 ds \quad (72)$$

$$= \frac{CN\kappa\rho}{(T-Ct)^2} \dot{y}\left(\frac{r}{T-Ct}\right) + \frac{4\pi\rho}{r^2} \int_0^r \frac{e^{y(s/(T-Ct))}}{(T-Ct)^3} s^2 ds \quad (73)$$

$$= \frac{\rho}{(T-Ct)^2} \left[CN\kappa \dot{y}\left(\frac{r}{T-Ct}\right) + \frac{4\pi}{r^2(T-Ct)} \int_0^r e^{y(s/(T-Ct))} s^2 ds \right]. \quad (74)$$

By letting $\omega = s/(T-Ct)$, it follows:

$$= \frac{\rho}{(T-Ct)^2} \left[CN\kappa \dot{y}\left(\frac{r}{T-Ct}\right) + \frac{4\pi}{\left(\frac{r}{T-Ct}\right)^2} \int_0^{r/(T-Ct)} e^{y(s)} s^2 ds \right] \quad (75)$$

$$= \frac{\rho}{(T-Ct)^2} Q\left(\frac{r}{T-Ct}\right). \quad (76)$$

And denote $z = r/(T - Ct)$,

$$Q\left(\frac{r}{T - Ct}\right) = Q(z) = CN\kappa\dot{y}(z) + \frac{4\pi}{z^2} \int_0^z e^{y(\omega)} \omega^2 d\omega. \quad (77)$$

Differentiate $Q(z)$ with respect to z ,

$$\dot{Q}(z) = CN\kappa\ddot{y}(z) + 4\pi e^{y(z)} + \frac{(-2) \cdot 4\pi}{z^3} \int_0^z e^{y(\omega)} \omega^2 d\omega \quad (78)$$

$$= -\frac{2CN\kappa}{z} \dot{y}(z) + \frac{(-2)4\pi}{z^3} \int_0^z e^{y(\omega)} \omega^2 d\omega \quad (79)$$

$$= -\frac{2}{z} Q(z), \quad (80)$$

where the above result is due to the fact that we choose the Liouville equation:

$$\begin{cases} \ddot{y}(z) + \frac{2}{z} \dot{y}(z) + \frac{4\pi}{CN\kappa} e^{y(z)} = 0 \\ y(0) = \alpha, \dot{y}(0) = 0. \end{cases} \quad (81)$$

With $Q(0) = 0$, this implies that $Q(z) = 0$. Now we are able to show that the family of the solutions blows up, in the finite time T/C . This completes the proof. ■

The statement about the blowup rate will be immediately followed:

Corollary 4 *The blowup rate of the solution (36) and (37) is*

$$\lim_{t \rightarrow \frac{T}{C}^-} \rho(t, 0)(T - Ct)^N \geq O(1). \quad (82)$$

Remark 5 *Besides, if we consider the Navier-Stokes equations with the repulsive force in radial symmetry in R^N ($N \neq 1$),*

$$\begin{cases} \rho_t + V\rho_r + \rho V_r + \frac{N-1}{r} \rho V = 0, \\ \rho(V_t + VV_r) = +\frac{\alpha(N)\rho}{r^{N-1}} \int_0^r \rho(t, s) s^{N-1} ds + [\kappa\rho^\theta]_r \left(\frac{N-1}{r} V + V_r\right) + (\kappa\rho^\theta)(V_{rr} + \frac{N-1}{r} V_r - \frac{N-1}{r^2} V) \end{cases} \quad (83)$$

the special solutions are:

for $N \geq 2$ and $N \neq 3$, with $\theta = (2N - 3)/N$,

$$\begin{cases} \rho(t, r) = \begin{cases} \frac{1}{(T - Ct)^N} y\left(\frac{r}{T - Ct}\right)^{\frac{N}{N-3}}, & \text{for } \frac{r}{T - Ct} < Z_0; \\ 0, & \text{for } Z_0 \leq \frac{r}{T - Ct} \end{cases}, \quad V(t, r) = \frac{-C}{T - Ct} r; \\ \ddot{y}(z) + \frac{N-1}{z} \dot{y}(z) - \frac{\alpha(N)(N-3)}{(2N-3)CN\kappa} y(z)^{\frac{N}{N-3}} = 0, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0, \end{cases} \quad (84)$$

the finite Z_0 is the first zero of $y(z)$;

for $N = 3$, with $\theta = 1$,

$$\begin{cases} \rho(t, r) = \frac{1}{(T - Ct)^3} e^{y(r/(T - Ct))}, \quad V(t, r) = \frac{-C}{T - Ct} r; \\ \ddot{y}(z) + \frac{2}{z} \dot{y}(z) - \frac{4\pi}{CN\kappa} e^{y(z)} = 0, \quad y(0) = \alpha > 0, \quad \dot{y}(0) = 0. \end{cases} \quad (85)$$

References

- [1] M. Bezard, *Existence locale de solutions pour les equations d'Euler-Poisson. (French) [Local Existence of Solutions for Euler-Poisson Equations]* Japan J. Indust. Appl. Math. **10** (1993), no. 3, 431–450.
- [2] S. Chandrasekhar, *An Introduction to the Study of Stellar Structure*, Univ. of Chicago Press, 1939.
- [3] Y.B. Deng, J.L. Xiang and T. Yang, *Blowup Phenomena of Solutions to Euler-Poisson Equations*, J. Math. Anal. Appl. **286** (1)(2003), 295-306.
- [4] P. Gamblin, *Solution reguliere a temps petit pour l'equation d'Euler-Poisson. (French) [Small-time Regular Solution for the Euler-Poisson Equation]* Comm. Partial Differential Equations **18** (1993), no. 5-6, 731–745.
- [5] P. Goldreich and S. Weber, *Homologously Collapsing Stellar Cores*, Astrophys. J. **238**, 991 (1980).
- [6] R. Kippenhahn, A. Weigert, *Stellar Structure and Evolution*, Springer-Verlag, 1990.
- [7] L.D. Landau and E.M. Lifshitz, E. M. (1959). *Fluid mechanics*. Translated from the Russian by J. B. Sykes and W. H. Reid. Course of Theoretical Physics. Vol. 6. London: Pergamon Press.
- [8] T. Makino, *Blowing up Solutions of the Euler-Poisson Equation for the Evolution of the Gaseous Stars*, Transport Theory and Statistical Physics **21** (1992), 615-624.
- [9] A. Mellet and A. Vasseur, *On the barotropic compressible Navier-Stokes equations*, Comm. Partial Differential Equations **32** (2007), no. 1-3, 431–452.
- [10] T. Nishida, *Equations of fluid dynamics-free surface problems*, Comm. Pure Appl. Math., XXXIX (1986), 221–238.
- [11] J. S. W., Wong, *On the generalized Emden-Fowler equation*, SIAM Rev. **17** (1975), 339-360.
- [12] T. Yang, Z.A. Yao and C. J. Zhu, *Compressible Navier-Stokes equations with density-dependent viscosity and vacuum*, Comm. Partial Differential Equations **26** (2001), no. 5-6, 965–981.
- [13] M.W. Yuen, *Blowup Solutions for a Class of Fluid Dynamical Equations in R^N* , J. Math. Anal. Appl. **329** (2)(2007), 1064-1079.
- [14] M.W. Yuen, *Analytical Blowup Solutions to the 2-dimensional Isothermal Euler-Poisson Equations of Gaseous Stars*, J. Math. Anal. Appl. **341** (1)(2008), 445-456.

- [15] M.W. Yuen, *Analytical Solutions to the Navier-Stokes Equations*, J. Math. Phys., **49** (2008) No. 11, 113102, 10pp.